



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae

Citation for published version:

Armstrong, K, Stone, G, Nicholls, J, Valderrama Escallon, E, Anderberg, A, Smedmark, J, Gautier, L, Naciri, Y, Milne, R & Richardson, JE 2014, 'Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae', *Frontiers in genetics*, vol. 5, 362. <https://doi.org/10.3389/fgene.2014.00362>

Digital Object Identifier (DOI):

[10.3389/fgene.2014.00362](https://doi.org/10.3389/fgene.2014.00362)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Frontiers in genetics

Publisher Rights Statement:

© 2014 Armstrong, Stone, Nicholls, Valderrama Escallon, Anderberg, Smedmark, Gautier, Naciri, Milne and Richardson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae

Kate Armstrong, Graham Stone, James Nicholls, Eugenio Valderrama_Escallon, Arne Anderberg, Jenny Smedmark, Laurent Gautier, Yamama Naciri, Richard Milne and James Edward Richardson

Journal Name: Frontiers in Genetics

ISSN: 1664-8021

Article type: Original Research Article

Received on: 18 Jul 2014

Accepted on: 29 Sep 2014

Provisional PDF published on: 29 Sep 2014

www.frontiersin.org: www.frontiersin.org

Citation: Armstrong K, Stone G, Nicholls J, Valderrama_escallon E, Anderberg A, Smedmark J, Gautier L, Naciri Y, Milne R and Richardson JE(2014) Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae. *Front. Genet.* 5:362. doi:10.3389/fgene.2014.00362

Copyright statement: © 2014 Armstrong, Stone, Nicholls, Valderrama_escallon, Anderberg, Smedmark, Gautier, Naciri, Milne and Richardson. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](http://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after rigorous peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae

K.E. Armstrong^{*,1,2,3}, G.N. Stone², J.A. Nicholls², E. Valderrama Escallón^{2,3}, A.A. Anderberg⁴, J. Smedmark⁵, L. Gautier⁶, Y. Naciri⁶, R. Milne⁷ & J.E. Richardson^{3,8}

¹ The New York Botanical Garden, Bronx, New York, U.S.A.

² Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, Scotland.

³ Royal Botanic Garden Edinburgh, Edinburgh, Scotland.

⁴ Naturhistoriska Riksmuseet, Stockholm, Sweden.

⁵ University Museum of Bergen, Bergen, Norway.

⁶ Conservatoire et Jardin botaniques, Genève, Switzerland.

⁷ Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh, Scotland.

⁸ Laboratorio de Botánica y Sistemática, Universidad de los Andes, Bogotá DC, Colombia.

Correspondence:

Kate Armstrong

The New York Botanical Garden

2900 Southern Boulevard

Bronx, New York, 10458, U.S.A.

karmstrong@nybg.org

Abstract

Species diversity is unequally distributed across the globe, with the greatest concentration occurring in the tropics. Even within the tropics, there are significant differences in the numbers of taxa found in each continental region. *Manilkara* is a pantropical genus of trees in the Sapotaceae comprising *c.* 78 species. Its distribution allows for biogeographic investigation and testing of whether rates of diversification differ amongst tropical regions. The age and geographical origin of *Manilkara* are inferred to determine whether Gondwanan break-up, boreotropical migration or long distance dispersal have shaped its current disjunct distribution. Diversification rates through time are also analyzed to determine whether the timing and tempo of speciation on each continent coincides with geoclimatic events. Bayesian analyses of nuclear (ITS) and plastid (*rpl32-trnL*, *rps16-trnK* and *trnS-trnFM*) sequences were used to reconstruct a species level phylogeny of *Manilkara* and related genera in the tribe Mimosoideae. Analyses of the nuclear data using a fossil-calibrated relaxed molecular clock indicate that *Manilkara* evolved 32-29 million years ago (Mya) in Africa. Lineages within the genus dispersed to the Neotropics

26-18 Mya and to Asia 28-15 Mya. Higher speciation rates are found in the Neotropical *Manilkara* clade than in either African or Asian clades. Dating of regional diversification correlates with known palaeoclimatic events. In South America, the divergence between Atlantic coastal forest and Amazonian clades coincides with the formation of drier Cerrado and Caatinga habitats between them. In Africa diversification coincides with Tertiary cycles of aridification and uplift of the east African plateaux. In Southeast Asia dispersal may have been limited by the relatively recent emergence of land in New Guinea and islands further east *c.* 10 Mya.

Key words

Sapotaceae, *Manilkara*, pantropical, biogeography, diversification rates

Introduction

Biodiversity is unevenly distributed across the globe and is most intensely concentrated in the tropics, particularly in wet tropical forests, which are the most species-rich biomes on the planet. Even within the tropics, there are significant differences in the floristic composition and the numbers of taxa found in each of the continental regions. It is estimated that there are *c.* 27,000 species of flowering plants in tropical Africa (Lebrun 2001; Lebrun & Stork 2003), compared with *c.* 90,000 for South America (Thomas 1999) and *c.* 50,000 for Southeast Asia (Whitmore 1998). This uneven species diversity raises the fundamental question of how variation in the pattern and tempo of speciation and extinction among continents might have driven observed patterns. Differences in diversity have been attributed to higher extinction rates in Africa (Richards 1973) and faster diversification in the Neotropics (Gentry 1982). Dated molecular phylogenies suggest speciation in response to recent climatic changes (such as aridification, e.g. Simon *et al* 2009, Couvreur *et al* 2008) or geological phenomena (such as mountain uplift in the Neotropics, e.g. Richardson *et al* 2001, Hughes & Eastwood 2006).

Intercontinental disjunctions in distribution between tropical regions of Africa, Asia and South America have been attributed to Gondwanan break-up (Raven & Axelrod 1974), and/or the degradation of the boreotropical flora (e.g. Malpighiaceae, Davis *et al* 2002; Meliaceae, Muellner *et al* 2006; Moraceae, Zerega *et al* 2006). However, current studies have shown that many tropical groups are of more recent origin (e.g. *Begonia*, Thomas *et al*, 2012), and that long distance dispersal has been an important factor in determining the composition of modern tropical floras (Pennington *et al*, 2006; Christenhusz & Chase, 2012). While long-distance dispersal could have occurred at any time, it was generally believed to be the only viable explanation for tropical intercontinental disjunctions younger than *c.* 33 Mya (although see Zhou *et al* 2012).

Pantropically distributed taxa are excellent models for studying the evolution of tropical forests and regional variation in diversification rates between continents. *Manilkara* is a genus of trees in the Sapotaceae comprising *c.* 78 species distributed throughout the tropics (30 in South and Central America, 35 in Africa and 13 in Southeast Asia). This even spread and relatively low number of species across major global tropical regions

93 makes *Manilkara* an excellent candidate for comparison of regional diversification
94 patterns and testing of hypotheses for the genesis of pantropical distributions. Here a near
95 species-level dated phylogeny of *Manilkara* is presented. If the distribution of the genus
96 can be explained by Gondwanan break up, the timing of phylogenetic splits would be
97 expected to reflect that break up 165-70 Mya (McLoughlin 2001). Similarly if splits
98 resulted from the degradation of the boreotropical flora, they would be expected to occur
99 as temperatures cooled following the Early Eocene Climatic Optimum/Paleocene–Eocene
100 Thermal Maximum (EECO/PETM), 50-55 Mya (Zachos 2001). Additionally, a
101 boreotropical origin should leave a phylogeographic signature in the form of southern
102 lineages being nested within more northern ones. Therefore, lineages in South America or
103 to the east of Wallace’s Line would be nested within Laurasian lineages, resulting in the
104 pattern one would expect from a retreat of the boreotropical flora from the Northern
105 Hemisphere. The onset of glaciation from 33 Mya induced further global cooling (Zachos
106 et al 2001) and the disintegration of the boreotropical flora. Therefore, ages of splits
107 younger than c. 33 Mya would most likely be explained by long distance dispersal. The
108 prediction advanced by Gentry (1983) that diversification rates in the Neotropics have
109 been higher than in other tropical regions is also tested.

111 **Materials and methods**

113 **DNA extraction, PCR, sequencing and alignment**

115 Evolutionary relationships were reconstructed using nuclear (ITS) and plastid (*rpl32-*
116 *trnL*, *rps16-trnK* and *trnS-trnFM*) sequences. Divergence times were calculated using an
117 ITS dataset with 171 accessions of Sapotaceae. In total 53 of the global total of 79
118 *Manilkara* species (67%) were included in the analysis. The dataset includes
119 representatives of the tribe Mimosopeae as well as multiple representatives of the tribes
120 Isonandreae and Sideroxyleae, which also belong to the subfamily Sapotoideae, in order
121 to accommodate calibration of fossils related to those groups. The tree was rooted using
122 *Sarcosperma*, shown in previous studies to be sister to the rest of the family (Anderberg
123 & Swenson 2003). The plastid dataset comprised 95 accessions of subtribe Manilkarinae,
124 as well as outgroups in subtribe Mimosopinae, plus *Northia*, *Inhambanella*, *Eberhardtia*
125 and *Sarcosperma*, which provided the root for the tree. See Supplementary Table 1 for
126 the list of taxa with voucher specimen information and GenBank accession numbers.

128 Total DNA was extracted from herbarium specimens and silica gel-dried leaf samples
129 using the Qiagen Plant DNeasy Mini Kit following the manufacturer’s instructions.
130 Amplifications of the ITS region were performed using the ITS5p/ITS8p/ITS2g/ITS3p
131 (Moeller & Cronk, 1997) and ITS1/ITS4 (White et al 1990) primer pairs. Polymerase
132 chain reaction (PCR) was carried out in 25-µL volume reactions containing 1 µL of
133 genomic DNA, 5.75 µL sterile distilled water, 2.5 µL 2 mM dNTPs, 2.5 µL 10x NH₄
134 reaction buffer, 1.25 µL 25 mM MgCl₂, 0.75 µL of each 10µM primer, 10 µL 5M betaine,
135 0.25 µL BSA and 0.25 µL of 5u/µL Biotaq DNA polymerase buffer. The thermal cycling
136 profile consisted of five minutes denaturation at 95°C, followed by 35 cycles of 30
137 seconds at 95°C for denaturation, 50°C for 30 seconds for annealing and 72°C for 1
138 minute and 30 seconds for extension with a final extension period of eight minutes at

72°C on a Tetrad2 BioRad DNA Engine. Extraction from herbarium specimens often yielded low amounts of degraded DNA and required nested PCR to amplify quantities sufficient for sequencing. In nested PCR we first used the ITS5/ITS8 primer pair, from which 1 µl of the PCR product was used in a second PCR with the ITS1/ITS4 primer pair and the same thermocycling profile. Further internal primers, ITS2g and ITS3p, were used in place of ITS1 and ITS4 when amplification using the latter primers was unsuccessful. Plastid markers were amplified using *rpl32-trnL* (Shaw et al 2007), *rps16-trnK* (Shaw et al 2007), and *trnS-trnFM* (Demesure et al 1995) primer pairs as well as *Manilkara*-specific internal primers designed for this study (Supplementary Table 2). PCR was carried out in 25 µL volume reactions containing 1 µL of genomic DNA, 15.25 µL sterile distilled water, 2.5 µL 2 mM dNTPs, 2.5 µL 10x NH₄ reaction buffer, 1.25 µL 25 mM MgCl₂, 0.75 µL of each 10 µM primer, 0.8 µL BSA and 0.2 µL of 5u/µL Biotaq DNA polymerase buffer. All plastid regions were amplified using the *rpl16* program of Shaw *et al* (2005). Nested PCR was also performed on selected accessions using self-designed internal primers (Supplementary Table 2). PCR products were purified using Exo-SAP (GE Healthcare) according to the manufacturer's instructions.

Sequencing PCRs were carried out using the BigDye Terminator v. 3.1 Cycle Sequencing Kit (Applied Biosystems) and were purified and sequenced on an ABI 3730 sequencer at the University of Edinburgh's GenePool facility. Forward and reverse sequences were assembled into contiguous sequences (contigs) and edited using the alignment software Sequencher ver. 4.7. Edited contigs were assembled and aligned by eye in MacClade ver. 4.08 (Maddison & Maddison 2008) and later in BioEdit ver. 7.0.5 (Hall 2005).

Potentially informative indels in the plastid dataset were coded according to the simple indel coding method of Simmons & Ochoterena (2000). Ambiguous alignment regions 113-118 and 380-459 in *rps16-trnK* were excluded. Indel events in ITS were so frequent that their coding as additional characters was deemed to be too ambiguous. Gaps were treated as missing data and all characters were equally weighted.

The ITS dataset was partitioned into three segments: ITS1 (372 bp), 5.8s (167 bp) and ITS2 (339 bp). Plastid regions and their indels were retained as separate partitions: *rpl32-trnL* (1130 bp + 26 indels), *rps16-trnK* (1134 bp + 21 indels) and *trnS-trnFM* (999 bp + 13 indels).

Phylogenetic analysis

Bayesian analyses were carried out using MrBayes 3.1 (Huelsenbeck & Ronquist 2001). Two independent runs of four MCMCMC chains each (three heated and one cold) were run with a temperature setting of 0.10 for 8,000,000 generations, which was found to provide sufficient mixing between chains and convergence between runs. Trees were sampled every 8,000 generations and a 10% burn-in was removed from the sampled set of trees leaving a final sample of 900 trees, which were used to produce a majority rule consensus tree. Convergence of models was determined to have occurred when the standard deviation of split frequencies for two runs reached 0.01 (Ronquist *et al* 2005). Appropriate burn-in and model convergence were checked by visual confirmation of

parameter convergence of traces in Tracer v.1.5 (Rambaut & Drummond 2009). Clade support values are posterior probabilities (pp); pp values of 100-95% are taken to indicate strong support, values of 94-90% moderate support, and values between 89-55% weak support for nodes, respectively. The output tree files were visualized in FigTree v.1.3.1. The majority rule consensus tree was used to determine the monophyly of key clades used to define calibration points in the dating analysis.

Plastid data were not included in the subsequent BEAST analysis because they were not informative enough to discern between alternative hypotheses and because fewer taxa were sampled. Additionally, hard incongruence was demonstrated between the topologies reconstructed in MrBayes from the nuclear and plastid datasets (see supplementary information section on chloroplast capture, and Figure S1). Therefore, the two datasets were not combined and only nuclear data was used for divergence time analysis.

Fossil calibration

Sideroxyleae pollen from the Ypresian (47.8-56 Mya) of England (Gruas-Cavagnetto, 1976) was used to constrain the minimum age of the Sideroxyleae stem node (node B in Fig.1). A log normal prior was used to constrain the age of this node (offset: 52.2 Ma, mean: 0.001). A mean of 0.001 was chosen so that 95% of the probability is contained in an interval between the midpoint and the upper boundary of the Ypresian (52.2–55.6 Mya). A Mid-Eocene (37.2-48.6 Mya) *Tetracolporpollenites* pollen grain from the Isle of Wight was used to constrain the minimum age of the node for the tribe Mimosopeae. This pollen grain was described by Harley (1991) and determined to closely resemble *Tieghemella heckelii* (a monotypic genus in the Mimosopeae). Harley suggested (pers. comm. 2010) that it would be appropriate to err on the side of caution with the identification and use the fossil to constrain the age of the tribe Mimosopeae rather than the genus itself. This fossil was, therefore, used to constrain the age of the crown node of Mimosopeae (node D in Fig.1: offset: 42.9 Mya, mean: 0.095). A mean of 0.095 was chosen so that 95% of the probability was contained in an interval between the midpoint (42.9) and the upper boundary of the mid Eocene (42.9-48.6 Mya). The final calibration point is based on a series of Oligocene (23-33.9 Mya) fossil leaves from Ethiopia (Jacobs *et al.*, 2005). Pan described these specimens as *Sapoteae* sp. and suggested possible placement in either *Manilkara* or *Tieghemella* (pers. comm. 2010) based on the occurrence of stoma surrounded by fimbriate periclinal rings, a character present in these genera, but absent from the related genera *Autranella* and *Mimusops*. Although they are both members of the Tribe Mimosopeae, *Manilkara* and *Tieghemella* are not sister taxa, and placing the fossil at the node of the most recent common ancestor (the entire Tribe Mimosopeae) seemed illogical for such a young date, when a 45 Mya fossil pollen grain of cf. *Tieghemella* was a better fit for the same node. Instead, the fossil was alternatively placed at the *Manilkara* crown node (node Q in Fig.1) and on the node of the split between *Tieghemella* and *Autranella* (node I in Fig.1), in order to determine whether placement on either genus made a significant difference to age estimates using a prior age estimate with an offset of 28 Mya, mean: 0.1. A mean of 0.1 was chosen so that 95% of the probability was contained in an interval between the midpoint and the upper boundary of the Oligocene at (28-33.9 Mya).

Dating analysis

The software package BEAST v.1.7.5 (Drummond & Rambaut 2007) was used to analyze divergence times in the ITS dataset. An xml input file was created in BEAUti v.1.7.5. Substitution models were unlinked across partitions, but clock models and tree topologies were kept on the linked default setting. Four taxon sets per analysis were generated in order to define nodes for placement of fossil calibration points. They were based on known monophyletic clades from previous analyses and were constrained to be monophyletic.

The GTR + I + G model was applied to each partition. The mean substitution rate was not fixed and base frequencies were estimated. Following support for a molecular clock in these data using MrBayes, an uncorrelated log-normal model was selected to allow for relaxed clock rates and rate heterogeneity between lineages. A speciation: birth-death process tree prior was used with a randomly generated starting tree. The most recent common ancestor (MRCA) node age priors were set to define calibration points using taxon sets. All other priors were left at default settings that were either uniform or gamma-distributed. Posterior distributions for each parameter were estimated using a Metropolis Coupled Monte Carlo Markov Chain (MCMCMC) run for 40,000,000 generations, with parameters logged every 5,000 generations, giving 8,000 samples per run. The BEAUti xml file was executed in BEAST v.1.7.5. Two separate analyses were run and the output log files were reviewed in Tracer v.1.5 (Rambaut & Drummond 2009) to check for convergence between runs and adequate effective sampling sizes (ESS) of > 200 (Drummond *et al* 2007). The tree files from the two runs were combined in LogCombiner v.1.7.5 (Drummond & Rambaut 2007) with a conservative burn-in of 4,000 generations. The combined tree files were input into TreeAnnotator v.1.5.3 (Drummond & Rambaut 2007). The Maximum Clade Credibility (MCC) tree was selected with mean node heights; this option summarizes the tree node height statistics from the posterior sample with the maximum sum of posterior probabilities. The output file was visualised in FigTree v.1.3.1.

Ancestral area reconstruction in RASP

Ancestral area states were reconstructed in RASP (Reconstruct Ancestral State in Phylogenies; <http://mnh.scu.edu.cn/soft/blog/RASP>) software that implements Bayesian Binary MCMC (BBM) time-events curve analysis (Yu et al., 2011) and allows multiple states to be assigned to terminals. BBM suggests possible ancestral ranges at each node and also calculates probabilities of each ancestral range at nodes. The analysis was performed using the MCC tree generated in BEAST as an input file, with 5,000,000 cycles, ten chains, sampling every 100 cycles, with a temperature setting of 0.1 and with the maximum number of areas set to four for all nodes. The root node was defined *a priori* as Asian; because the Asian taxa *Sarcosperma* and *Eberhardtia* form a grade within which the rest of the family is nested, this is the most likely state for the crown node of the family.

Areas are coded according to continent, based predominantly on tectonic plate margins and then on floristic regions (Fig. 1). In Southeast Asia, the Sahul and Sunda Shelves (which mark the boundary between continental Asia and Australia-New Guinea) were coded as separate states within the Malesia floristic region, which stretches from the Isthmus of Kra on the Malay Peninsula to Fiji. East Asia is defined as being east of the Himalayas and south as far as the Malay Peninsula, with a predominantly Indo-Chinese flora. South Asia is delineated by the margin of the Indian subcontinent. The countries of Iran, Turkey and the Arabian Peninsula support a drier Irano-Turanian flora and were, therefore, designated as being part of the Middle-Eastern region. The remaining regions (the Seychelles, Madagascar, Africa and North and South America) are all on separate continental tectonic plates and are floristically unique from one another (see Supplementary Table 1 for species-specific area codes).

Diversification rate methods

A separate ITS lineage through time plot dataset (hereafter referred to as ITS LTT) was used to compare diversification rates within *Manilkara*. Because the genus was found to be paraphyletic, with the Southeast Asian *M. fasciculata* clade (P in Fig. 1) being more closely related to *Labourdonnaisia* and *Faucherea*, this small clade was excluded, leaving only the monophyletic lineage of *Manilkara* s.s. (clade Q in Fig. 1) for analysis. Additionally, only one individual per species was included. The simple diversification rate estimators of Kendall (1949) and Moran (1951) were calculated for the African, Neotropical and Asian clades, where the speciation rate $SR_{ln} = [\ln(N) - \ln(N_0)]/T$ (N = standing diversity, N_0 = initial diversity, here taken as = 1, and T = inferred clade age). This is a pure-birth model of diversification with a constant rate and no extinction (Magallón and Sanderson, 2001). Another model that does not assume constant rates of speciation and extinction through time within lineages was applied using BAMM (Bayesian Analysis of Macroevolutionary Mixtures; Rabosky 2014). BAMM uses a reversible-jump Markov Chain Monte Carlo to explore shifts between macroevolutionary regimes, assuming they occur across the branches of a phylogenetic tree under a compound Poisson process. Each regime consists in a time-varying speciation rate (modeled with an exponential change function) and a constant rate of extinction. The BAMM analysis used the BEAST MCC tree, but because not all species were sampled, it was necessary to specify to which lineage each of the missing taxa belonged, (i.e. to which species it was most closely related based on morphological similarity). Two MCMC simulations were run with 5,000,000 generations, sampling every 1,000, and discarding the first 10% as burn-in. Appropriate priors for the ITS LTT phylogeny, convergence of the runs and effective sampling size were each estimated using the BAMM tools package in R (Rabosky 2014).

Lineage through time (LTT) plots were generated using phytools (Revell 2012) in R (R development team) for 1000 trees sampled through the post-burn-in (20%) posterior distribution generated by BEAST (see above for details). The median and 95% highest posterior density (HPD) were estimated for the ages of each number of lineages in each plot. To compare the observed LTT plots with the predictions of a model with constant diversification rates, 1000 trees were simulated using the mean speciation and extinction

rates estimated by BAMM in TreeSim (Stadler 2011). Simulations used the age of the most recent common ancestor of each of the 1000 observed trees and the current number of species per plot. LTT plots were drawn for the trees including all species of *Manilkara* s.s. and to examine region-specific patterns for pruned lineages that included only those species from each of Africa, the Neotropics and Asia.

Results

Node ages

Mean ages with 95% HPD confidence intervals for key nodes are reported in Table 1. The MCC tree from the BEAST analysis (Fig.1) resolves the mean crown age of the tribe Mimosopeae as 43 Mya (HPD 44-42 Mya; node D), in the Mid Eocene. The mean age of subtribe Manilkarinae is estimated to be 32 Mya (HPD 36-29 Mya; node K) and the genus *Manilkara* is resolved as 29 Mya (HPD 32-28 Mya; node Q), both having originated during the Oligocene. Results also reveal that cladogenesis and inter-continental dispersal (see below and Fig.1, Fig.3) within *Manilkara* occurred from the Oligocene through the Miocene – and most intensively from the mid-late Miocene.

Table 1. Summary of clade support values, node ages and ancestral areas from Figure 1.

Node	Posterior probability	Clade	Mean age and 95% HPD in Mya	Ancestral Area (likelihood %)	Epoch
A	1	Sapotaceae	107 (126-88)	East Asia 99%	Cretaceous
B	1	Sideroxyleae	62 (73-52)	Africa 58%	Cretaceous-Paleocene
C	0.99	Isonandreae/ <i>Inhambanella</i> /Mimosopeae	52 (58-48)	Africa 99%	Paleocene-Eocene
D	1	Mimosopeae	43(44-42)	Africa 99%	Eocene
E	0.99	<i>Baillonella</i> / <i>Vitellaria</i> / <i>Vitellariopsis</i>	31(39-23)	Africa 99%	Eocene-Oligocene
F	0.99	<i>Vitellariopsis</i>	2 (4-0.5)	Africa 99%	Pliocene
G	0.85	Mimosopeae subclade 1	39 (43-35)	Africa 99%	Eocene
H	0.67	<i>Mimusops</i> / <i>Tieghemella</i> / <i>Autranella</i>	35 (40-30)	Africa 99%	Eocene-Oligocene
I	0.68	<i>Tieghemella</i> / <i>Autranella</i>	31 (38-23)	Africa 99%	Eocene-Oligocene
J	0.99	<i>Mimusops</i>	22 (28-17)	Africa 97%	Miocene
K	0.99	Manilkarinae	32 (36-29)	Africa 96%	Eocene-Oligocene
L	0.44	<i>Labr./Fauch./Labourdonnaisia</i> /sm. Asian <i>Manilkara</i>	30 (35-26)	Madagascar 81%	Eocene-Oligocene
M	0.99	<i>Labramia</i>	6 (10-3)	Madagascar 99%	Miocene-Pliocene
N	0.92	<i>Faucherea</i> / <i>Labourdonnaisia</i> / <i>Manilkara</i>	28 (33-23)	Madagascar 91%	Oligocene
O	0.99	<i>Faucherea</i> / <i>Labourdonnaisia</i>	10 (14-7)	Madagascar 99%	Miocene-Pliocene
P	0.99	Small Asian <i>Manilkara</i>	15 (20-10)	Sahul shelf 90%	Miocene
Q	1	<i>Manilkara</i> s.s.	29 (32-28)	Africa 96%	Oligocene
R	0.98	<i>Manilkara</i> s.s. subclade 1	26 (30-22)	Africa 86%	Oligocene-Miocene
S	0.99	Neotropical <i>Manilkara</i>	18 (22-14)	South America 71%	Miocene
T	0.90	Central American & Caribbean <i>Manilkara</i>	15 (20-13)	North America 95%	Miocene
U	0.99	South American <i>Manilkara</i> s.s.	12 (16-9)	South America 93%	Miocene
V	0.77	Small African <i>Manilkara</i>	21 (27-15)	Africa 97%	Oligocene
W	0.99	<i>Manilkara</i> s.s. subclade 2	27(30-23)	Africa 97%	Oligocene
X	0.99	Large African <i>Manilkara</i>	15 (18-11)	Africa 99%	Miocene
Y	0.99	Asian <i>Manilkara</i> s.s.	23 (27-19)	Sahul Shelf 52%	Oligocene-Miocene

Ancestral area reconstruction and intercontinental dispersal events

Ancestral area inferences and likelihood support are given in Table 1 and Figure 1, which also indicates the age and direction of inferred dispersal events. The tribe Mimosopeae,

subtribe *Manilkarinae* and the genera *Manilkara*, *Labramia* and *Faucherea/Labourdonnaisia* are all inferred to have African ancestry (Fig. 1).

Following its origin in Africa during the Oligocene 32 Mya (HPD 36-29; node K) and subsequent diversification 29 Mya (HPD 32-28 Mya; node Q), *Manilkara s.s.* spread via long distance dispersal to Madagascar twice, Asia once and the Neotropics once during the Oligocene–Miocene. Both the *Faucherea/Labourdonnaisia/ Manilkara* clade (N) (28 Mya; HPD 33-23 Mya) and the genus *Mimusops* (clade J) (22 Mya; HPD 28-17 Mya) also exhibit a similar pattern, having originated in Africa and later dispersed to both Madagascar and Asia during the Miocene.

Long-distance dispersal from Africa to Madagascar and the surrounding islands has occurred on multiple occasions in the tribe *Mimosopeae*: twice in *Manilkara s.s.* (X3 & X4, 8-4 Mya); at least once for the clade comprising *Labramia*, *Faucherea*, and *Labourdonnaisia* between 32 Mya (HPD 36-29; node K) and 30 Mya (HPD 35-26 Mya; node L); and twice in *Mimusops* between 22 Mya (HPD 28-17 Mya; node J) and 9 Mya (HPD 13-5 Mya; node J1), as well as 5 Mya (HPD 2 – 6 Mya; node J3).

The Neotropical *Manilkara* clade (S) is also derived from an African ancestor, which dispersed to South America during the Oligocene–Miocene between 26 Mya (HPD 30-22 Mya; node R) and 18 Mya (HPD 22-14 Mya; node S). From South America, further dispersal occurred to Central America 16-15 Mya and throughout the Caribbean islands starting from 15-10 Mya.

Asia was reached by three independent dispersal events within the tribe *Mimosopeae*. *Manilkara s.s.* reached Asia from Africa between 27 Mya (HPD 30-23 Mya; node W) and 23 Mya (HPD 27-19 Mya; node Y), while *Mimusops* did the same 8-6 Mya (node J2). The *Manilkara fasciculata* clade reached Asia from Madagascar between 28 (HPD 33-23 Mya; node N) and 15 Mya (HPD 20-10 Mya; node P).

Diversification rates

Net diversification rates (SRln) differed somewhat between regions, ranging from a lowest mean value of 0.06 (0.05-0.07) for the Asian lineage, through 0.10 (0.09-0.10) for the African lineage to a maximum of 0.15 (0.12-0.19) for the Neotropical lineage. Despite sampling models with up to five different macroevolutionary regimes, BAMM analysis consistently selected models without shifts between macroevolutionary regimes along the *Manilkara* phylogeny, with the highest posterior probability obtained for zero shifts models, i.e. a single, constantly varying net diversification rate throughout the history of the genus (Figure 2).

Lineage through time (LTT) plots are presented in Figure 3, for all regions (Fig. 3d) and for the pruned African, Asian and Neotropical lineages (Fig. 3a-c respectively). The figure shows both observed rates, and rates predicted for the same numbers of lineages evolving under a constant net diversification rate process (i.e. constant speciation and extinction rates, estimated using BAMM for the whole genus). None of the observed LTT

patterns diverge significantly from those predicted assuming a constant diversification rate. The analyses including all *Manilkara* lineages (Fig. 3d) and only the Neotropical lineage (Fig. 3c) both show a good fit between observed patterns and those predicted under a constant diversification rate. In contrast, African lineages (Fig. 3a) show a trend towards reduced diversification rates from 25 to 12 Mya, followed by an increase in diversification rates to levels matching those in the Neotropics from 12 Mya to the present. The Asian lineage shows low and decreasing diversification rates towards the present. While the Asian pattern is derived from just six species, and thus any observed pattern must be interpreted with caution, it is striking that Asia produced no new lineages during the last 7 Mya, at a time when Africa and the Neotropics were both showing rapid diversification.

Discussion

Origin of *Manilkara*

The tribe Mimosopeae evolved approximately 52 Mya (HPD 58-48 Mya; node C) and began to diversify 43 Mya (HPD 44-32 Mya; node D) during the Eocene when global climates were warmer and wetter and a megathermal flora occupied the northern hemisphere. This age estimate also coincides with the first occurrence of putative Mimosopeae fossils recorded from North America and Europe, e.g. *Tetracolporpollenites brevis* (Taylor 1989), and *Manilkara* pollen (Frederiksen 1980) in addition to the *Tetracolporpollenites* sp., pollen grain (Harley 1991), used in this study, which give further weight to the hypothesis that the tribe Mimosopeae was present in the boreotropics and may have originated there. Previous studies (Smedmark & Anderberg 2007) implicate the break-up of the boreotropics in creating intercontinental disjunctions in the tribe Sideroxyleae and data from the present study are consistent with this hypothesis. Smedmark & Anderberg's (2007) estimate for the age of Sideroxyleae was 68 Mya and in this study the crown node age is reconstructed as being 62 Mya (HPD 73-52 Mya; node B).

The subtribe Manilkarinae evolved 39 Mya (HPD 43-35 Mya; node G), consistent with the hypothesis that it arose late during the existence of the boreotropics. Diversification began 32 Mya (HPD 36-29 Mya; node K), around the time that global cooling and the widening Atlantic were breaking up the boreotropics. Hence migration towards the equator as the climate in the northern hemisphere cooled might have caused or promoted diversification. This transition from the northern hemisphere to equatorial latitudes is also reflected in the putative Manilkarinae fossil record, where during the Oligocene, there is still a strong representation of northern fossils (e.g. Isle of Wight, U.K. (Machin 1971), Vermont, U.S.A. (Traverse 1953 & 1955) and Czechoslovakia (Prakash, Brezinova & Awasthi 1974)), but fossils also begin to appear in Africa (e.g. *Sapoteae* sp. leaves in Ethiopia (Jacobs *et al* 2005)). Further cooling and aridification during the Oligocene coincides with diversification of Manilkarinae into genera and may have been a causal factor in this diversification. Alternatively, Manilkarinae may have originated in Africa, as suggested by the ancestral area analysis. However, the analysis cannot account for southward climate shifts and the modern absence of the group from higher latitudes.

Manilkara is nested within a grade of other representatives of the tribe Mimosopeae, which is predominantly composed of African taxa (*Mimusops*, *Tieghemella*, *Autranella*, *Baillonella*, *Vitellaria* and *Vitellariopsis*) and this suggests that the genus may have had its origin there. In the ancestral area reconstruction both *Manilkara* and the subtribe Manilkarinae are resolved as having a 96% likelihood of an African origin, and the tribe Mimosopeae is reconstructed as having a 99% likelihood of originating in Africa. As such, there is very strong support for an African ancestry for the genus *Manilkara*, the subtribe Manilkarinae and the tribe Mimosopeae.

The origin of *Manilkara*'s pantropical distribution

Intercontinental disjunctions in *Manilkara* are too young (27-4 Mya) to have been caused by Gondwanan break-up, which would have had to occur before 70 Mya. *Manilkara* is also too young for its pantropical distribution to be the result of migration through the boreotropics, which would have had to occur between 65-45 Mya, after which the climate would have been too cool for tropical taxa to cross the North Atlantic Land Bridge, even though this might have persisted until ~33 MYA (Milne and Abbott, 2002). The most likely period for migration of tropical taxa by this route was during the PETM/EECO, 55-50 Mya (Zachos 2001). Furthermore, a boreotropical origin should leave a phylogeographic signature in the form of southern lineages being nested within more northern ones. However, South American lineages are not nested within Central American lineages, and neither are those southeast of Wallace's line nested within those to the northwest. With these vicariance-based explanations not supported, *Manilkara*'s disjunct pantropical distribution could only have resulted from long-distance dispersal from Africa to Madagascar, Asia and the Neotropics. This has been demonstrated for numerous other groups distributed across the tropics, e.g. *Begonia* (Thomas et al 2012) and *Renealmia* (Sarkinen et al, 2007).

Manilkara has fleshy, sweet fruit ranging in size from 1.5 – 10cm, which are consumed by a wide variety of animals. With seeds that are too bulky for wind dispersion, it is more likely that long distance dispersal could have been achieved through transport in the gut-contents of birds or by transoceanic rafting in large mats of vegetation. Houle's (1998) study demonstrated that during the Miocene, intercontinental rafting could have occurred in less than two weeks on the North and South Equatorial currents.

Regional diversification in *Manilkara*

Within the Neotropics, *Manilkara* first colonized South America, as indicated in the reconstruction of the ancestral distribution of clade S. The South American clade (U) is divided into two subclades, which correspond to contrasting regional ecologies, with one clade (U1) comprised of Amazonian species and the other (U2) of Atlantic coastal forest species. The only inconsistency in this geographic pattern is the second accession of *Manilkara cavalcantei* (b), an Amazonian species that the analysis places in the Atlantic coastal forest clade. However, in the plastid tree (Supplementary Figure 1) this accession is resolved in a strongly supported (0.99 pp) Amazonian clade with *M. bidentata*, *M.*

487 *huberi* and *M. paraensis*. The phylogenetic split between these two regions occurred
488 during the Mid-Miocene (12-10 Mya), when the Andes were being elevated (Graham
489 2009; Gregory-Wodzicki 2000) and drainage systems in the Amazon basin began to shift
490 eastwards.

491
492 Atlantic coastal species in clade U2 and Amazonian species in clade U1 are
493 geographically separated by the dry biomes of the Cerrado and the Caatinga, as well as
494 the higher relief of the Brazilian shield. Simon *et al* (2009) and Fritsch *et al* (2004) found
495 that the origin of dry-adapted Cerrado Leguminosae and Melastomataceae lineages span
496 the Late Miocene to the Pliocene (from 9.8 to 0.4 Mya), broadly coinciding with the
497 expansion of C4 grass-dominated savanna biomes. However, it is likely that a dry
498 environment would have been present just prior to this time to allow for adaptation of
499 these groups to the new biome. Such timing is exhibited by the Microlicieae
500 (Melastomataceae), where the crown node is 9.8 Mya, and the stem node is 17 Mya
501 (Fritsch *et al* 2004). *Manihot* (Euphorbiaceae) species of this biome began to diversify
502 from 6.6 Mya (Chacon *et al*, 2008). Likewise, a phylogenetic study of *Coursetia*
503 (Leguminosae) (Lavin 2006) reveals that species which inhabit the dry forest of the
504 Brazilian Caatinga are 5-10 My old. This suggests that the Cerrado and Caatinga could
505 have been in existence, at least in part, by the time the South American *Manilkara*
506 subclades U1 & U2 diverged ca.12 Mya, and their development may have driven the
507 geographical split in this South American lineage of *Manilkara*.

508
509 African *Manilkara* species are resolved in two clades, both of which are Oligo-Miocene
510 in age. The main African/Madagascan clade (X) is estimated to be 15 My old (HPD 18-
511 11 Mya), and the smaller clade (V) is 21 My old (HPD 27-15 Mya). Africa has been
512 affected by widespread aridification during the Tertiary (Coetzee 1993, Morley 2000).
513 The response by *Manilkara* to this changing climate could have been migration,
514 adaptation or extinction. A study of the rain forest genera *Isolona* and *Monodora*
515 (Annonaceae) found that throughout climatic cycles, taxa remained in remnant pockets of
516 wet forest (Couvreur *et al* 2008). They are, therefore, an example of a group that
517 migrated or changed its distribution to track wetter climates. Another study of the genus
518 *Acridocarpus* (Malpighiaceae) (Davis *et al* 2002) indicated an east African dry forest
519 adapted lineage nested within a wet forest lineage. The dry adapted lineage was dated to
520 periods of Oligo-Miocene aridification, and is, therefore, an example of a wet forest
521 lineage, which has adapted to changing environmental conditions rather than becoming
522 restricted to areas of favorable climate. The timing of diversification and evolution of
523 dry-adapted species versus wet-restricted species in the three African *Manilkara* clades
524 suggests a combination of both scenarios. The split between the African clades occurred
525 between 29 Mya (HPD 32-28 Mya; node Q) and 26 Mya (HPD 30-22 Mya; node R),
526 during a period of dramatic continent-wide cooling, which fragmented the Eocene coast
527 to coast rain forest, potentially isolating the three lineages. A second wave of
528 diversification within the main African/Madagascan clade (X) coincides with the Mid-
529 Miocene climatic optimum 17-15 Mya, when global temperatures warmed (Zachos
530 2001). During the same period the collision of the African and Eurasian plates closed the
531 Tethys Sea, instigating further aridification. The resulting drier and warmer climates
532 caused the spread of savannas and the retraction of rain forest, as evidenced by an

increase in grass pollen during this period (Morley 2000; Jacobs 2004). Nonetheless, cladogenesis in the main African/Madagascan clade (X) gained pace from the Mid-Miocene onwards. In particular, a third wave of diversification from rain forest into drier shrubland environments in eastern and southern Africa occurred subsequent to the main uplift of the Tanganyikan plateau in the East African Rift System ca. 10 Mya, which had a significant impact on further regional aridification (Lovett & Wasser 1993; Sepulchre *et al* 2006) (Table 1).

Clade X is predominantly composed of Guineo-Congolian rain forest species. This is almost exclusively the case in subclade X1, aside from the Madagascan taxa, which are also rain forest species. However, within subclade X2, there is a transition from wet to dry environments. The sole Madagascan taxon in this lineage (*M. sahafarensis*) is a dry, deciduous forest species. The four dry, eastern-southern African taxa in subclade X2 (*M. discolor*, *M. sansibarensis*, *M. butugi*, *M. cuneifolia*) all evolved between 8-5 Mya subsequent to the main uplift of the East African Rift System. The ancestor of the smaller African clade composed of *M. mochisia* and *M. concolor* also diversified into these two dry-adapted eastern/southern species at the same time 6 Mya (HPD 10-2 Mya). Hence, some African *Manilkara* lineages adapted to a drying climate, while others remained in their ancestral rain forest habitat.

Within the main Asian clade of the plastid phylogeny (Yc1, Supplementary Figure 1), the Indian species *Manilkara roxburghiana* is sister to the other species and the two Fijian species are among the most derived, consistent with the hypothesis that the founding dispersal event was from Africa to India with subsequent spread eastward into Malesia. However, ancestral area reconstruction of the ITS data (node Y, Figure 1) suggests that migration within Asia was from east to west (Sahul Shelf to Sunda Shelf) 23 Mya (HPD 27-19 Mya). Dated phylogenies also indicate that many other angiosperm groups have crossed Wallace's Line from the late Miocene onwards: *Pseuduvaria* (Annonaceae) (Su & Saunders 2009), Aglaieae (Meliaceae) (Muellner *et al* 2008), at least four separate lineages of *Begonia* (Begoniaceae) (Thomas *et al* 2012) and *Cyrtandra* (Gesneriaceae) (Cronk *et al* 2005). In Sapotaceae four lineages of Isonandreae have migrated from west to east across Wallace's Line (Richardson *et al* 2014), whereas evidence from the tribe Chrysophylloideae suggests recent movement in the opposite direction, from Sahul to Sunda Shelf (Swenson *et al* 2013). The two youngest (9 Mya) Asian species (*M. vitiensis* & *M. smithiana*) are both Fijian. The oldest land available for colonization in Fiji is between 14-5 Mya (Johnson 1991; Heads 2006) hence, the age of these two Fijian taxa coincides with the first emergence of land in the archipelago.

Diversification rates of *Manilkara* in different parts of the tropics

The BAMM analysis did not support significant rate variation among lineages or regions in *Manilkara s.s.* Despite apparent variation in regional patterns revealed by lineage through time plots (Fig.3), the data most strongly support a model with a single net diversification rate throughout the genus. Trends within the data for specific regions only suggest departure from a constant rate model in Asia and Africa. Given that observed patterns do not exceed the 95% confidence intervals for the constant rate model for either

region, these trends must be considered with caution. This is particularly true for Asia, for which the pattern was derived from only eight species. Because sensitivity and statistical power of methods for detection of shifts in diversification rates may correlate positively with the number of species in the clade (Silvestro, 2012), rate shifts in clades with a small number of species (as in Asia for *Manilkara s.s.*) may not have been detected by the methods used here (a potential type two error). A simulation study would be required to examine the impact of taxon number on type two error rates in these analyses. Similarly, small numbers of taxa may be more likely to generate apparent trends through stochastic effects, and these could also generate the apparent two-phase pattern of low, and then rapid, diversification in African lineages.

Taken at face value, net diversification rates and LTT plots both suggest a trend for more rapid diversification in Neotropical and African lineages than in Asian ones. The timing of rapid Neotropical diversification falls within the time frame of Andean uplift (i.e. from the late Miocene onwards), proposed as a diversification engine in many taxa (e.g. Richardson et al 2001). However, because many South American *Manilkara* species are native to the Atlantic Forest, on the opposite side of the continent from the Andes, Andean uplift may be considered unlikely to directly explain high diversification rates region-wide. Interestingly, the rapid diversification of the African lineage coincided with periods of regional aridification. The slowest diversification rate, in the Southeast Asian lineage, includes species that are mostly to the east of Wallace's Line. This may be explained by the fact that the mountainous topography of much of this region (dominated by New Guinea) limits the habitat available for lineages such as *Manilkara* that are largely restricted to lowland rain forest that covers a greater area of Africa or the Neotropics. Although there is no statistical support for significant diversification rate variation in *Manilkara s.s.*, the causes highlighted here should have similar impacts on other lowland rainforest taxa – a predication that can be tested in future studies utilizing phylogenies of more species rich taxa and meta-analyses of multiple unrelated lineages.

Author Contributions

This paper is a result of KA's Ph.D. thesis research at the Royal Botanic Garden Edinburgh and University of Edinburgh. KA and JER conceived the study and KA carried out the research and wrote the manuscript apart from the diversification rate analysis, which was conducted and written by EVE. JER, GS and RM supervised the Ph.D. project. GS and JER edited the manuscript. JN assisted with phylogenetic analyses. AAA, JS, LG and YN contributed DNA sequence data to the study. All authors have reviewed the manuscript.

Acknowledgements

This doctoral research was made possible through a scholarship from the Torrance Bequest at the University of Edinburgh. Grants for fieldwork from the Royal Geographical Society, the Carnegie Trust, the Systematics Association, and the Davis Expedition Fund are also gratefully acknowledged. D. Ndiade Bourobou (CENAREST, IRAF) is thanked for a DNA aliquot of *Baillonella toxisperma*, and Jerome Chave

(CNRS) is thanked for ITS sequences of *Manilkara bidentata* and *M. huberi*. Thanks to members of the Stone Lab at the University of Edinburgh for comments on an earlier draft of the manuscript.

References

Anderberg, A.A. & Swenson, U. (2003). Evolutionary lineages in Sapotaceae (Ericales): A cladistic analysis based on *ndhF* sequence data. *Int. J. Plant Sci.* 164(5): 763-773.

Burgess, N. D., Clarke, G. P. and Rodgers, W. A. (1998). Coastal forests of eastern Africa: status, endemism patterns and their potential causes. *Biol. J. Linn. Soc.* 64(3): 337-367. doi: <http://dx.doi.org/10.1111/j.1095-8312.1998.tb00337.x>

Chacon Pinilla, J., Madriñán, S., Tohme, J., Ebouck, D., and Rodriguez, F. (2008). Phylogenetic patterns in the genus *Manihot* (Euphorbiaceae) inferred from analyses of nuclear and chloroplast DNA regions. *Mol. Phylogenet. Evol.* 49(1): 260-7. doi: 10.1016/j.ympev.2008.07.015.

Chan, K., and B. R. Moore. (2005). SYMMETREE: Whole-tree analysis of differential diversification rates. *Bioinformatics.* 21(8): 1709-1710. doi: 10.1093/bioinformatics/bti175.

Christenhusz, M. J. M. and Chase, M. W. (2013). Biogeographical patterns of plants in the Neotropics – dispersal rather than plate tectonics is most explanatory. *Bot. J. Linn. Soc.* 171(1): 277–286. doi: 10.1111/j.1095-8339.2012.01301.x.

Coates, A. G. and Obando, J. A. (1996). “The geologic evolution of the Central American isthmus,” in *Evolution and environment in tropical America*, eds. J. B. C. Jackson, A. F. Budd, & A. G. Coates (Chicago, IL: University of Chicago Press), 21-56.

Cody, S., Richardson, J. E., Rull, V., Ellis, C. and Pennington, R. T. (2010). The great American biotic interchange revisited. *Ecography* 33: 1-7. doi: 10.1111/j.1600-0587.2010.06327.

Coetzee, J. A. (1993). “African Flora since the terminal Jurassic,” in *Biological Relationships between Africa and South America*, ed. P. Goldblatt (New Haven, CT: Yale University Press), 37-61.

Couvreur, T. L. P., Chatrou, L. W., Sosef, M. S. and Richardson, J. E. (2008). Molecular phylogenetics reveal multiple tertiary vicariance origins of the African rain forest trees. *BMC Biol.* 6:54. doi:10.1186/1741-7007-6-54.

Crisp, M., Cook, L. and Steane, D. (2004). Radiation of the Australian flora: what can comparisons of molecular phylogenies across multiple taxa tell us about the evolution of diversity in present-day communities? *Philos. Trans. R. Soc. Lond., B.* 359(1450): 1551-1571. doi: 10.1098/rstb.2004.1528

- Cronk, Q. C. B., Kiehn, M., Wagner, W. L. and Smith, J. F. (2005). Evolution of *Cyrtandra* (Gesneriaceae) in the Pacific Ocean: the origin of a supertramp clade. *Am. J. Bot.* 92(6): 1017-1024. doi: 10.3732/ajb.92.6.1017
- Davis, C. C., Bell, C. D., Fritsch, P. W. and Matthews, S. (2002a). Phylogeny of *Acridocarpus-Brachylophon* (Malpighiaceae): implications for Tertiary tropical floras and Afroasian biogeography. *Evolution* 56(12): 2395-2405. doi: [http://dx.doi.org/10.1554/0014-3820\(2002\)056\[2395:POABMI\]2.0.CO;2](http://dx.doi.org/10.1554/0014-3820(2002)056[2395:POABMI]2.0.CO;2)
- Davis, C. C., Bell, C. D., Matthews, S. and Donoghue, M. J. (2002b). Laurasian migration explains Gondwanan disjunctions: evidence from Malpighiaceae. *Proc. Natnl. Acad. Sci. USA* 99(10): 6833-6837. doi: 10.1073/pnas.102175899.
- Demesure, B., Sodzi, N. and Petit, R. J. (1995). A set of universal primers for amplification of polymorphic noncoding regions of mitochondrial and chloroplast DNA in plants. *Mol. Ecol.* 4(1): 129-131.
- Drummond, A. J. and Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* 7: 214. doi:10.1186/1471-2148-7-214.
- Drummond, A. J., Ho, S. Y. W., Rawlence, N. and Rambaut, A. (2007). Beast Manual: A Rough Guide to BEAST 1.4. Available at: <http://beast.bio.ed.ac.uk>.
- Erkens, R.H. J., Maas, J. W. and Couvreur T.L.P. (2009). From Africa via Europe to South America: migrational route of a species-rich genus of Neotropical lowland rainforest trees (*Guatteria*, Annonaceae). *J. Biogeogr.* 36(12): 2338-2352. doi: 10.1111/j.1365-2699.2009.02162.x
- Farris, D.W., Jaramillo, C., Bayona, G., Restrepo-Moreno, S.A., Montes, C. Cardona, A., Mora, A., Speakman, R.J., Glascock, M.D., Valencia, V. (2011). Fracturing of the Panamanian Isthmus during initial collision with South America. *Geology*. 39(11): 1007-1010. doi: 10.1130/G32237.1
- Frederiksen, N. O. (1980). Mid-Tertiary climate of Southeastern United-States - The sporomorph evidence. *J. Paleontol.* 54(4): 728-739.
- Fritsch, P. W., Almeda, F., Renner, S. S., Martins, A. B. and Cruz, B. C. (2004). Phylogeny and circumscription of the near-endemic Brazilian tribe Microlicieae (Melastomataceae). *Am. J. Bot.* 91(7): 1105-1114. doi: 10.3732/ajb.91.7.1105.
- Gentry A.H. (1982). Floristic diversity: phytogeographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny? *Ann. Mo. Bot. Gard.* 69, 557-593. doi: <http://dx.doi.org/10.2307/2399084>.

- Graham, A. (2009). The Andes: a geological overview from a biological perspective. *Ann. Mo. Bot. Gard.* 96(3): 371-385. doi: <http://dx.doi.org/10.3417/2007146>.
- Gregory-Wodzicki, K. M. (2000). Uplift history of the Central and Northern Andes: A review. *Geol. Soc. Am. Bull.* 112(7): 1091-1105. doi: 10.1130/0016-7606(2000)112<1091:UHOTCA>2.0.CO;2.
- Gruas-Cavagnetto, C. (1976). Etude palynologique du Paleogene du sud de l'Angleterre. *Cah. Micropal.* 1: 1-49.
- Haffer, J. (1969). Speciation in Amazonian forest birds. *Science* 165(3889): 131-137. doi: 10.1126/science.165.3889.131
- Hall, R. (1998). "The plate tectonics of Cenozoic SE Asia and the distribution of land and sea," in *Biogeography and geological evolution of SE Asia*. eds R. Hall and J.D. Holloway (Leiden, The Netherlands: Backhuys Publishers) 99-131.
- Hall, R. (2001). "Cenozoic reconstructions of SE Asia and the SW Pacific: Changing patterns of land and sea," in *Faunal and Floral Migrations and Evolution in SE Asia-Australasia*. eds I. Metcalfe, J.M.B. Smith, M. Moorwood and I.D. Davidson (Lisse, The Netherlands: A.A. Balkema; Swets & Zeitlinger Publishers). 35-56.
- Hall, R. (2002). Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. *J. Asian Earth Sci.* 20(4): 353-431. doi: [http://dx.doi.org/10.1016/S1367-9120\(01\)00069-4](http://dx.doi.org/10.1016/S1367-9120(01)00069-4).
- Hall, R. (2009). Southeast Asia's changing palaeogeography. *Blumea* 54(1-3): 148-161. doi: <http://dx.doi.org/10.3767/000651909X475941>.
- Hall, T. (2005). BioEdit version 7.0.5. Carlsbad, California: Ibis Therapeutics. Available at: <http://www.mbio.ncsu.edu/bioedit/bioedit.html>.
- Harley, M. M. (1991). The pollen morphology of the Sapotaceae. *Kew Bull.* 46(3): 379. doi: <http://dx.doi.org/10.2307/4110538>.
- Heads, M. (2006). Seed plants of Fiji: an ecological analysis. *Biol. J. Linn. Soc.* 89(3): 407-431. doi: 10.1111/j.1095-8312.2006.00682.x.
- Houle, A. (1998). Floating islands: a mode of long distance dispersal for small and medium-sized terrestrial vertebrates. *Divers. Distrib.* 4: 201-216.
- Huelsenbeck, J. P. and Ronquist, F. (2001). MrBayes. Bayesian inference of phylogenetic trees. *Bioinformatics* 17(8): 754-755. doi: <http://dx.doi.org/10.1093/bioinformatics/17.8.754>

- Hughes, C. and Eastwood, R. (2006). Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the Andes. *Proc. Natnl. Acad. Sci. USA*. 103(27): 10334-10339. doi: 10.1073/pnas.0601928103.
- Jacobs, B. F. (2004). Palaeobotanical studies from tropical Africa: relevance to the evolution of forest, woodland and savannah biomes. *Phil. Trans. R. Soc. Lond. B* 359: 1573-1583. doi: 10.1098/rstb.2004.1533.
- Jacobs, B. F., Tabor, N., Feseha, M., Pan, A., Kappelman, J., Rasmussen, T., Sanders, W., Wiemann, M. C., Crabaugh, J. and Garcia Massini, J. L. (2005). Oligocene terrestrial strata of northwestern Ethiopia: a preliminary report on paleoenvironments and paleontology. *Palaeontologica Electronica* 8(1): 1-19. http://palaeo-electronica.org/paleo/2005_1/jacobs25/issue1_05.htm.
- Johnson, H. (1991). Petroleum geology of Fiji. *Mar. Geol.* 98: 2-4. doi: [http://dx.doi.org/10.1016/0025-3227\(91\)90109-H](http://dx.doi.org/10.1016/0025-3227(91)90109-H)
- Kass, R. E. and Raferty, A. E. (1995). Bayes Factors. *J. Am. Stat. Assoc.* 90(430): 773-795. doi: <http://dx.doi.org/10.2307/2291091>
- Kendall, D. G. (1949). Stochastic processes and population growth. *J. R. Stat. Soc. B Stat. Methodol.* 11, 230-264.
- Lavin, M., Schrire, B., Lewis, G., Pennington, T.R., Delgado-Salinas, A., Thulin, M., Hughes, C., Matos, A.B., and Wojciechowski, M.F. (2004). Metacommunity process rather than continental tectonic history better explains geographically structured phylogenies in legumes. *Philos. Trans. R. Soc. Lond., B*. 359: 1509-1522. doi:10.1098/rstb.2004.1536.
- Lavin, M. (2006). "Floristic and geographical stability of discontinuous seasonally dry tropical forests explains patterns of plant phylogeny and endemism," in (eds.) *Neotropical Savannas and Seasonally Dry Forests: Plant Diversity, Biogeography, and Conservation*, eds. R.T. Pennington, G.P. Lewis and J.A. Ratter (Boca Raton, FL: CRC Press) 433-447.
- Lebrun, J. P. and Stork, A. L. (2003). *Tropical African flowering plants: ecology and distribution, volume 1 Annonaceae-Balanitaceae*. Conservatoire et Jardin Botaniques de la Ville de Geneve.
- Lebrun, J. P. (2001). *Introduction à la flore d'Afrique*. Paris: Cirad, Ibis Press.
- Lovett, J. and Wasser, S. (1993). *Biogeography and Ecology of the Rainforests of Eastern Africa*. Cambridge: Cambridge University Press.
- Machin, J. (1971). Plant microfossils from Tertiary deposits of the Isle of Wight. *New Phytol.* 70: 851-872. doi: <http://dx.doi.org/10.1111/j.1469-8137.1971.tb02586.x>.

- Maddison, W. P. and Maddison, D. R. (2008). MacClade version 4.08. Sinauer Associates, Sunderland. Available at: <http://macclade.org/macclade.html>.
- Magallón, S. and Sanderson, M. J. (2001). Absolute diversification rates in angiosperm clades. *Evolution* 55, 1762–1780. doi: <http://dx.doi.org/10.1111/j.0014-3820.2001.tb00826.x>.
- Manen, J. F. Barriera, G., Loizeau, P. A., and Naciri, Y. (2010). The history of extant *Ilex* species (Aquifoliaceae): Evidence of hybridization within a Miocene radiation. *Mol. Phylogenet. Evol.* 57: 961–2977. doi: 10.1016/j.ympev.2010.09.006.
- McLoughlin, S. (2001). The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. *Australian Journal of Botany*. 49(3): 271-300.
- Milne, R. I. and Abbott, R. J. (2002). The origin and evolution of tertiary relict floras. *Adv. Bot. Res.* 38: 281-314. doi: [http://dx.doi.org/10.1016/S0065-2296\(02\)38033-9](http://dx.doi.org/10.1016/S0065-2296(02)38033-9).
- Möller, M. and Cronk, Q. C. B. (1997). Origin and relationships of *Saintpaulia* (Gesneriaceae) based on ribosomal DNA internal transcribed spacer (ITS) sequences. *Am. J. Bot.* 84(7): 956-965. doi: <http://dx.doi.org/10.2307/2446286>.
- Moran, P. A. (1951). Estimation methods for evolutive processes. *J. R. Stat. Soc. B Stat. Methodol.* 13, 141–146.
- Morley, R. J. (2000). *Origin and evolution of tropical rain forests*. Chichester: John Wiley & Sons, Ltd.
- Muellner, A. N., Pannell, C. M., Coleman, A. and Chase, M. W. (2008). The origin and evolution of Indomalayan, Australasian and Pacific island biotas: insights from Aglaieae (Meliaceae, Sapindales) *J. Biogeogr.* 35: 1769-1789. doi: <http://dx.doi.org/10.1111/j.1365-2699.2008.01935.x>.
- Muellner, A. N., Savolainen, V., Samuel, R. & Chase, M. W. 2006. The mahogany family “out-of-Africa”: divergence time estimation, global biogeographic patterns inferred from plastid *rbcL* DNA sequences, extant, and fossil distribution of diversity. *Mol. Phylogenet. Evol.* 40: 236–250.
- Nylander, J. A. A. (2008). MrModeltest version 2.3. Evolutionary Biology Centre, Uppsala University: Available at: <http://darwin.uvigo.es/software/modeltest.html>.
- Peng, D. and Wang, X. Q. (2008). Reticulate evolution in *Thuja* inferred from multiple gene sequences: Implications for the study of biogeographical disjunction between eastern Asia and North America. *Mol. Phylogenet. Evol.* 47(3): 1190-1202. doi: <http://dx.doi.org/10.1016/j.ympev.2008.02.001>.

- Pennington, R.T., Lavin, M., Prado, D.E., Pendry, C.A., Pell, S.K., Butterworth, C.A. (2004). Historical climate change and speciation: neotropical seasonally dry forest plants show patterns of both Tertiary and Quaternary diversification. *Philos. Trans. R. Soc. Lond., B.* 359(1443): 515-537. PMC1693336
- Pennington R.T., Richardson, J.E., and Lavin M. (2006). Insights into the historical construction of species-rich biomes from dated plant phylogenies, phylogenetic community structure and neutral ecological theory. *New Phytol.* 172(4): 605-616. doi: <http://dx.doi.org/10.1111/j.1469-8137.2006.01902.x>.
- Plana, V., Gascoigne, A., Forrest, L. L., Harris, D. J. and Pennington, T. R. (2004). Pleistocene and pre-Pleistocene *Begonia* speciation in Africa. *Mol. Phylogenet. Evol.* 31(2): 449–461. doi: <http://dx.doi.org/10.1016/j.ympev.2003.08.023>.
- Prakash, U., Brezinova, D. and Awasthi, N. (1974). Fossil woods from the Tertiary of South Bohemia. *Paleontographica Abteilung B* 147: 107-133.
- Rabosky, D.L. (2014). Automatic Detection of Key Innovations, Rate Shifts, and Diversity-Dependence on Phylogenetic Trees. *PLoS ONE* 9(2): e89543.
- Rambaut, A. and Drummond, A. J. (2009). Tracer v.1.5. Available at: <http://tree.bio.ed.ac.uk/software/tracer/>.
- Raven, P. H. and Axelrod, D. I. (1974). Angiosperm biogeography and past continental movements. *Ann. Mo. Bot. Gard.* 61(3): 539-673. doi: <http://dx.doi.org/10.2307/2395021>.
- Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217-223.
- Richards, P.W. 1973. "Africa, the Odd Man Out," in *Tropical forest ecosystems in Africa and South America: a comparative review*, eds. B. J. Meggers, E. S. Ayensu, and W D. Duckworth. (Washington: Smithsonian Institution Press) 21- 26.
- Richardson, J.E., Bakar, A.M., Tosh, J., Armstrong, K.E., Smedmark, J. Anderberg, A. Slik, F. and Wilkie, P. (2014). The influence of tectonics, sea-level changes and dispersal on migration and diversification of Isonandreae (Sapotaceae). *Bot. J. Linn. Soc.* 174(1) 130-140.
- Richardson, J. E., Pennington, R. T., Pennington, T. D. and Hollingsworth, P. M. (2001). Rapid diversification of a species-rich genus of neotropical rain forest trees. *Science* 293(5538): 2242-2245. doi: <http://dx.doi.org/10.1126/science.1061421>.
- Rodda, P. (1976). *Geology of northern and central Viti Levu*. Bulletin, Mineral Resources Division Fiji, volume 3. Suva, Fiji: Ministry of Lands and Mineral Resources.

- Rodda, P. (1994). "Geology of Fiji," in *Contributions to the marine and on-land geology and resources of the Tonga-Lau-Fiji region*, eds. A. J. Stevenson, R. H. Herzer and P. F. Ballance (Suva, Fiji: SOPAC Technical Bulletin) 131-151.
- Ronquist, F., Huelsenbeck, J. P. and van der Mark, P. (2005). MrBayes 3.1. Available at: <http://mrbayes.csit.fsu.edu/>.
- Sarkinen, T. E., Newman, M. F., Maas, P. J. M., Maas, H., Poulsen, A. D., Harris, D. J., Richardson, J. E., Clark, A., Hollingsworth, M. and Pennington, T. R. (2007). Recent oceanic long-distance dispersal and divergence in the amphi-Atlantic rain forest genus *Renealmia* L.f. (Zingiberaceae). *Mol. Phylogenet. Evol.* 44: 968-980. doi: <http://dx.doi.org/10.1016/j.ympev.2007.06.007>.
- Sarkinen, T., Pennington, R.T., Lavin, M., Simon, M.F., Hughes, C.E., (2012). Evolutionary islands in the Andes: persistence and isolation explain high endemism in Andean dry tropical forests. *J. Biogeogr.* 39(5): 884-900. doi: 10.1111/j.1365-2699.2011.02644.x
- Sepulchre, P., Ramstein, G., Fluteau, F., Schuster, M., Tiercelin, J. J. and Brunet, M. (2006). Tectonic uplift and Eastern Africa aridification. *Science* 313(5792): 1419-1423. doi: <http://dx.doi.org/10.1126/science.1129158>.
- Shaw, J., Lickey, E. B., Beck, J. T., Farmer, S. B., Liu, W. S., Miller, J., Siripun, K. C., Winder, C. T., Schilling, E. E. and Small, R. L. (2005). The tortoise and the hare II: Relative utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *Am. J. Bot.* 92(1): 142-166. doi: <http://dx.doi.org/10.3732/ajb.92.1.142>.
- Shaw, J., Lickey, E. B., Schilling, E. E. and Small, R. L. (2007). Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: The tortoise and the hare III. *Am. J. Bot.* 94(3): 275-288. doi: <http://dx.doi.org/10.3732/ajb.94.3.275>.
- Silvestro, D., Schnitzler, J. Zizka, G. 2011. A Bayesian framework to estimate diversification rates and their variation through time and space. *BMC Evolutionary Biology* 11:311.
- Simmons, M. P. and Ochoterena, H. (2000). Gaps as characters in sequence-based phylogenetic analyses. *Syst. Biol.* 49(2): 369-381. doi: <http://dx.doi.org/10.1093/sysbio/49.2.369>.
- Simon, M. F., Grether, R., de Queiroz, L. P., Skema, C., Pennington, R. T. and Hughes, C. E. (2009). Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire. *Proc. Natnl. Acad. Sci. USA.* 106(48): 20359-20364. doi: <http://dx.doi.org/10.1073/pnas.0903410106>.

- Smedmark, J. E. E. and Anderberg, A. A. (2007). Boreotropical migration explains hybridization between geographically distant lineages in the pantropical clade Sideroxyleae (Sapotaceae). *Am. J. Bot.* 94: 1491-1505. doi: <http://dx.doi.org/10.3732/ajb.94.9.1491>.
- Stadler, T. (2011). Simulating trees on a fixed number of extant species. *Syst. Biol.* 60: 676-684.
- Stebbins, G. L. (1974). *Flowering plants: evolution above the species level*. Cambridge, MA: Harvard University Press.
- Stone, G. N., Hernandez-Lopez, A., Nicholls, J. A., di Pierro, E., Pujade-Villar, J., Melika, G. & Cook, J. M. (2009). Extreme host plant conservatism during at least 20 million years of host plant pursuit by oak gallwasps. *Evolution*, 63, 854-869.
- Su, Y. C. F. and Saunders, R. M. K. (2009). Evolutionary divergence times in the Annonaceae: evidence of a late Miocene origin of *Pseuduvaria* in Sundaland with subsequent diversification in New Guinea. *BMC Evol. Biol.* 9(153). doi: <http://dx.doi.org/10.1186/1471-2148-9-153>.
- Swenson, U., Lowry, P. P., Munzinger, J., Rydin, C. and Bartish, I. V. (2008). Phylogeny and generic limits in the *Niemeyera* complex of New Caledonian Sapotaceae: evidence of multiple origins of the anisomerous flower. *Mol. Phylogenet. Evol.* 49(3): 909-929. doi: <http://dx.doi.org/10.1016/j.ympev.2008.09.022>.
- Taylor, D. W. (1989). Select palynomorphs from the Middle Eocene Caliborne Formation, Tenn, (USA). *Rev. Palaeobot. Palynol.* 58(2-4): 111-128.
- Thomas D.C., Hughes M., Phutthai T., Ardi W.H., Rajbhandary S., Rubite R., Twyford A.D., and Richardson J.E. (2012) West to east dispersal and subsequent rapid diversification of the mega-diverse genus *Begonia* (Begoniaceae) in the Malesian archipelago. *J. Biogeogr.* 39, 98–113. doi: <http://dx.doi.org/10.1111/j.1365-2699.2011.02596.x>.
- Thomas, W. W. (1999). Conservation and monographic research on the flora of tropical America. *Biodivers. Conserv.* 8: 1007–1015. doi: 10.1023/A:1008857429787.
- Traverse, A. (1955). *Pollen analysis of the Brandon Lignite of Vermont*. Report of Investigations 5151: United States Department of the Interior, Bureau of Mines.
- Traverse, A. and Barghoorn, E.S. (1953). Micropaleontology of the Brandon lignite, an early Tertiary coal in central Vermont. *J. Paleontol.* 27(2): 289-293.
- Verboom, G.A., Archibald, J.K., Bakker, F.T., Bellstedt, D.U., Conrad, F., Dreyer, L.L., Forest, F., Galley, C., Goldblatt, P., Hennig, J.F., Mummenhoff, K., Linder, H. P., Muasya, A.M., Oberlander, K.C., Savolainen, V., Snijman, D.A., van der Niet, T., and

- Nowell, T.I. (2009). Origin and diversification of the Greater Cape flora: ancient species repository, hot-bed of recent radiation, or both? *Mol. Phylogenet. Evol.* 51(1): 44-53. <http://dx.doi.org/10.1016/j.ympev.2008.01.037>
- Wendel, J. F., Schnabel, A. and Seelanan, T. (1995). An unusual ribosomal DNA sequence from *Gossypium gossypoides* reveals ancient, cryptic intergenomic introgression. *Mol. Phylogenet. Evol.* 4(3): 298-313. doi: <http://dx.doi.org/10.1006/mpev.1995.1027>.
- White, T. J., T. Burns, S. Lee, and J. Taylor. (1990). "Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics," in *PCR protocols: a guide to methods and applications* eds. M. Innis, D. Gelfand, J. Sninsky & T. White (San Diego, CA: Academic Press) 315-322.
- Whitmore, T. C. (1998). *An introduction to tropical rainforests*, 2nd ed. Oxford University Press.
- Yu Y., Harris A.J., and He X. (2011) RASP (Reconstruct Ancestral State in Phylogenies) 2.1.B, Available at: <http://mnh.scu.edu.cn/soft/blog/RASP>.
- Zachos, J. C., Pagani, M., Sloan, L., Thomas, E. and Billups, K. (2001). Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292: 686-693. doi: <http://dx.doi.org/10.1126/science.1059412>.
- Zerega, N. J. C., Clement, W. L., Datwyler, S. H. & Weiblen, G. D. 2005. Biogeography and divergence times in the mulberry family (Moraceae). *Molecular Phylogenetics and Evolution* 37: 402–416.
- Zhou, L., Su, Y.C.F., Thomas, D.C. and Saunders, R.M.K. 2012. 'Out-of-Africa' dispersal of tropical floras during the Miocene climatic optimum: evidence from Uvaria (Annonaceae). *Journal of Biogeography* 39: 322–335.

FIGURE LEGENDS

Figure 1. Maximum clade credibility chronogram of the ITS dataset. Dashed lines indicate branches which lead to nodes with a posterior probability of <0.95. Mean ages are given for profiled nodes. Node bars indicate 95% HPD age ranges. Lettered nodes are discussed in the text. Stars indicate the placement of fossils. Lineages are colored according to their distribution: Yellow = Africa, Green = Madagascar, Blue = Asia, Pink = South America, Orange = Central America & the Caribbean. Geological epochs are indicated in a scale at the bottom of the chronogram. Outgroups have been reduced to grey bars at the base of the chronogram. Ten regions were coded in the ancestral area reconstruction as illustrated in the map and legend. Pie charts represent the percentage likelihood of the ancestral state at the selected node. Map inset depicts the timing and direction of long-distance dispersal events reflected in the chronogram.

Figure 2. Posterior probability of models with different number of shifts between macroevolutionary regimes considered in BAMM. The best models for *Manilkara s.s.* indicate no significant shifts in diversification.

Figure 3. LTT plots for lineages that included only those species from each of Africa (a), Asia (b), the Neotropics (c) and all species of *Manilkara s.s.*(d). Each plot shows the median and 95% HPD of the ages for each number of lineages in solid and dashed lines, respectively. The lines for observed trees are shown in blue and for the trees simulated under a constant diversification process in red. The thinner blue lines correspond to each of the 1000 observed trees. The 95% HPD intervals show major overlap in all plots but non-significant patterns suggest lower diversification rates in part of the histories of African and Asian lineages.

SUPPLEMENTARY INFORMATION

Incongruence between nuclear and plastid trees

Phylogenies generated with nuclear (Fig. 1) and plastid data (Supplementary Fig. 1) showed high topological congruence. However, there are a couple examples of hard incongruence (strongly supported clades which conflict in their placement between the two datasets), both of which have biogeographic implications. The first is in the placement of the two Asian species *Manilkara hexandra* and *M. littoralis*, and the two African species *M. mochisia* and *M. concolor*. In the ITS phylogeny *M. hexandra* and *M. littoralis* are resolved in the Asian clade Y, while *M. mochisia* and *M. concolor* are resolved in the small African clade V. In contrast, in the plastid phylogeny, these four species form a strongly supported clade (posterior probability 1), marked Φ in Fig. S1.

A second hard incongruence is apparent in the placement of the three taxa *Manilkara yangambensis*, *M. triflora* and *M. suarezensis*. In the plastid phylogeny these form a monophyletic clade Z (Fig. S1). In contrast, in the ITS analysis, the Brazilian *M. triflora* was poorly resolved at the base of clade T, whereas the Madagascan *M. suarezensis* was resolved within the main African clade (X). The Congolese species *M. yangambensis* was not included in the ITS analysis due to difficulties in amplifying its DNA from herbarium specimens.

These discrepancies between the nuclear and plastid trees may be the result of either ancestral polymorphism with incomplete lineage sorting or chloroplast capture (introgression) following dispersal.

Hard incongruence between nuclear and plastid trees – evidence for chloroplast capture?

In the dated nuclear phylogeny, the Asian species *M. hexandra* (Sri Lanka) and *M. littoralis* (Myanmar) (clade Y1) are placed with other Asian species (clade Y2), whereas in the plastid phylogeny, they are resolved in clade Φ with two African species *M.*

mochisia (Zambia) and *M. concolor* (South Africa) (from clade V in ITS). This suggests hybridization of taxa across the Indian Ocean possibly resulting in chloroplast capture. Intercontinental chloroplast capture may also be implicated in the case of clade Z, which is resolved in the plastid analyses but not in the ITS analyses and is composed of *M. suarezensis* (Madagascar), *M. triflora* (Brazil) and *M. yangambensis* (Congo). The ITS analysis did not include *M. yangambensis*, but placed *M. triflora* with other Neotropical species in clade S, and *M. suarezensis* with other Madagascan species within a larger clade of African species (clade X). Therefore, ITS resolved at least two of the clade Z species with species from the same landmass, but cpDNA did not, and resolved them together instead. Clade Z is strongly supported (pp 0.99) in the plastid analysis. Assuming that the correct species level relationships are resolved, clade Z presents a case of long distance dispersal and chloroplast capture more remarkable than the clade V/Y1 scenario, because it involves species from three landmasses, and hence two dispersal events.

Hybridization and chloroplast capture across long distances such as ocean barriers has been indicated previously in Sapotaceae. The species *Chrysophyllum cuneifolium* is inferred to have originated from an intercontinental hybridization event where the chloroplast is South American and the nuclear genome is African (Swenson *et al* 2008). Likewise, the Pacific genus *Nesoluma* is hypothesized to have arisen as a result of intercontinental hybridization in the boreotropical region during the Eocene (Smedmark & Anderberg 2007). *Nesoluma* presents the opposite pattern to *Chrysophyllum*, where the chloroplast is African and the nuclear genome is Neotropical. Hybridization between New and Old World lineages has also been demonstrated in the pantropical genus *Gossypium* (Malvaceae) (Wendel *et al* 1995) and intercontinental chloroplast capture is hypothesized to have also occurred in *Thuja* (Cupressaceae) (Peng & Wang 2008). Additionally, both hybridization and introgression events are inferred to have occurred between distantly related species in *Ilex* (Aquifoliaceae) (Manen *et al* 2010). What is abundantly clear is that long distance dispersal has played a crucial role in the establishment of the modern distribution of *Manilkara*.

Supplementary Figure 1. Bayesian majority rule consensus tree of the chloroplast dataset. Posterior probability values are indicated above branches. Nodes with letters/symbols are discussed in the text.

Supplementary Table 1. Herbarium specimen data, GenBank accession number and ancestral area coding for taxa included in the analyses. Accessions of newly generated sequences are emboldened.

Supplementary Table 2. Chloroplast primers designed for this study.

Primer name	Direction	Primer sequence (5'-3')
rpl32-trnL-intF	forward	TCGTCGAGATTGAAGAGTCA
rpl32-trnL-intR	reverse	TCTCTTTTGACCGGAAATTCA
rpl32_trnL_int_2_F	forward	GGCGGCTGCTCAACTTAT
rpl32_trnL_int_2_R	reverse	TCTCTTTTGACCGGAAATTCA

rps16-trnK-intF	forward	TGTTCCCTGCTATTCTATATTTCCTTG
rps16-trnK-intR	reverse	GATGTGTAGATACAATCAGAATCAAAA
rps16_trnK_int_2_F	forward	GGGTGCTCAACCTACAGAAA
rps16_trnK_int_2_R	reverse	ACGAGGCAATCAAAACATTG
trnS-trnFM_int.F	forward	ACTCAGCCATCTCTCCGAAA
trnS-trnFM_int.R	reverse	TTTGGGGTGAGAGGAAAAGA
trnS-trnFM_int_2_F	forward	AACCACTCAGCCATCTCTCC
trnS-trnFM_int_2_R	reverse	GAACCCCTACACTATCACGG

1122

1123

Figure 1.JPEG




Figure 2.JPEG




Figure 3.JPEG

